


10/510019

DT04 Rec'd PCT/PTO 0 1 OCT 2004

**Amendments to the claims:**

The following listing of claims replaces all prior versions of claims in this application.

1. (Currently Amended) A method for the treatment of a disease or condition where an antagonist of the alpha-2 adrenoceptor is indicated to be useful, which comprises administering to a patient in need of the treatment an effective amount of Use of a compound of formula I,



wherein,

X is CR<sub>2</sub>R<sub>2</sub>', O, S or NR<sub>2</sub>;

Z is -CHR<sub>8</sub>-(CH<sub>2</sub>)<sub>n</sub>- or a single bond;

R<sub>1</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, CN, NO<sub>2</sub>, NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino or carboxyl;

R<sub>2</sub> and R<sub>2</sub>' are independently H, hydroxy or (C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>2</sub> and R<sub>2</sub>' form, together with the carbon ring atoms to which they are attached, a carbonyl group;

R<sub>3</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or one of R<sub>3</sub> or R<sub>4</sub> and R<sub>6</sub> together form a bond between the ring atoms to which they are attached;

R<sub>4</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or R<sub>4</sub> and R<sub>5</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring substituted with 1 to 3 substituent(s) R<sub>9</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-

C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>6</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>6</sub> forms a bond between the ring atom to which it is attached and the ring atom to which R<sub>7</sub> is attached;

R<sub>7</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>8</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or, only when n is 0, R<sub>7</sub> and R<sub>8</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring unsubstituted or substituted with 1 to 3 substituent(s) R<sub>10</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>15</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or carboxyl;

R<sub>16</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>7</sub> and R<sub>8</sub> are attached to the carbon ring atoms, which are adjacent;

m is 0 to 2; and

n is 0 or 1,

or a pharmaceutically acceptable salt or ester thereof, with the proviso, that the compound is not 1,2,3,4,5,10b-hexahydro-10-thia-3a-aza-cyclopenta[a]fluorine,

~~for the manufacture of a medicament for the treatment of diseases or conditions where antagonists of alpha-2 adrenoceptors are indicated to be useful.~~

2. (Currently Amended) A method ~~The use of a compound~~ according to claim 1, wherein X is NR<sub>2</sub>.

3. (Currently Amended) A method ~~The use of a compound~~ according to claim 1 ~~any one of claims 1 or 2~~, wherein m is 0, n is 0, R<sub>2</sub> is H, R<sub>3</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO- or (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sub>4</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl or hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, R<sub>5</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkoxy, R<sub>6</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sub>7</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl.

4. (Currently Amended) A method ~~The use of a compound~~ according to claim 1 ~~any one of claims 1 to 3~~, wherein R<sub>3</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sub>4</sub> is hydroxy or hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl.

5. (Currently Amended) A method ~~The use of a compound~~ according to claim 1 ~~any one of claims 1 or 2~~, wherein  $R_4$  and  $R_5$  form, together with the carbon ring atoms to which they are attached, a condensed six membered saturated carbocyclic ring.

6. (Currently Amended) A method ~~The use of a compound~~ according to claim 1 ~~one of claims 1 or 2~~, wherein  $R_4$  and  $R_6$  together form a bond between the ring atoms to which they are attached or  $R_6$  forms a bond between the ring atom to which it is attached and the ring atom to which  $R_7$  is attached.

7. (Currently Amended) A method ~~The use of a compound~~ according to claim 1 ~~any one of claims 1 to 5~~, wherein the compound is  $1\alpha$ -ethyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-a]quinolizin-1-ol, (1 $\beta$ -ethyl-1,2,3,4,6,7,12,12b $\alpha$ -octahydro-indolo[2,3-a]quinolizin-1-yl)-methanol,  $1\alpha$ -Methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-a]quinolizin-1-ol, (1 $\alpha$ -Methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-a]quinolizin-1-yl)-methanol or 3,4,4a $\beta$ , 5,6,7,8,13,13b $\beta$ ,13c $\alpha$ -decahydro-2H-6a,13-diaza-indeno[1,2-c]phenanthren-1-one.

8. (Currently Amended) A method ~~The use of a compound~~ according to claim 1, wherein X is  $CR_2R_2'$ .

9. (Currently Amended) A method ~~The use of a compound~~ according to claim 1, wherein X is O

10. (Currently Amended) A method ~~The use of a compound~~ according to claim 1, wherein X is S.

11. A method ~~The use of a compound~~ according to claim 1, which comprises any one of claims 1 to 10, for the manufacture of a medicament for the treatment of a disorder of


the central nervous system, diabetes, orthostatic hypotension, lipolytic disorder disorders, Raynaud's disease or male or and female sexual dysfunctions.

12. (Currently Amended) A method according to claim 11, wherein the disorder of the central nervous system is depression, anxiety disorder disorders, post-traumatic stress disorder, schizophrenia, Parkinson's disease, or another movement disorder.

13. (Currently Amended) A method ~~The use of a compound~~ according to claim 1, wherein the compound is any one of claims 1 to 10 for the manufacture of a medicament for use as a selective alpha-2C antagonist.

14. (Currently Amended) A method ~~The use~~ according to claim 13, which comprises for the manufacture of a medicament for the treatment of a mental disorder disorders propagated by stress, Parkinson's disease, depression, negative symptoms of schizophrenia, attention deficit hyperactivity disorder, post-traumatic stress-disorder, or anxiety disorder disorders.

15. (Original) A compound of formula IA



wherein,



C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or R<sub>4</sub> and R<sub>5</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring substituted with 1 to 3 substituent(s) R<sub>9</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>6</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>6</sub> forms a bond between the ring atom to which it is attached and the ring atom to which R<sub>7</sub> is attached;

R<sub>7</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>8</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or, only when n is 0, R<sub>7</sub> and R<sub>8</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring unsubstituted or substituted with 1 to 3 substituent(s) R<sub>10</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl,



carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>15</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or carboxyl;

R<sub>16</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>7</sub> and R<sub>8</sub> are attached to the carbon ring atoms, which are adjacent;

m is 0 to 2; and

n is 0 or 1,

or a pharmaceutically acceptable salt or ester thereof, with the provisos, that

a) when X is O, m is 0 and n is 0, then R<sub>3</sub>-R<sub>8</sub> are not all simultaneously hydrogen;


b) the compound is not 1,2,3,4,5,10b-hexahydro-10-thia-3a-aza-cyclopenta[a]fluorene; 1,3,4,5,6,11b-hexahydro-2H-11-thia-4a-aza-benzo[a]fluorene; 1-(1,3,4,5,6,11b-hexahydro-2H-11-thia-4a-aza-benzo[a]fluoren-1-yl)-ethanone or 1,3,4,5,6,11b-hexahydro-2H-11-thia-4a-aza-benzo[a]fluorene-1-carboxylic acid methyl ester.

16. (Original) A compound according to claim 15, wherein X is CR<sub>2</sub>R<sub>2</sub>'.

17. (Original) A compound according to claim 15, wherein X is O.
18. (Original) A compound according to claim 15, wherein X is S.
19. (Currently Amended) A compound according to claim 15 ~~any one of claims 15 to 18~~, wherein R<sub>3</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO- or (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sub>4</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl.
20. (Currently Amended) A compound according to claim 15 ~~any one of claims 15 to 19~~, wherein R<sub>3</sub> is hydroxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sub>4</sub> is (C<sub>1</sub>-C<sub>6</sub>)alkyl.
21. (Currently Amended) A compound according to claim 15 ~~any one of claims 15 to 18~~, wherein R<sub>4</sub> and R<sub>5</sub> form, together with the carbon ring atoms to which they are attached, a condensed six membered saturated carbocyclic ring.
22. (Currently Amended) A compound according to claim 15 ~~any one of claims 15 to 21~~, wherein the compound is 1 $\alpha$ -Methyl-1,3,4,5,6,11b-hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-ol, (1 $\alpha$ -Methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, (-)-(1 $\alpha$ -Methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, (+)-(1 $\alpha$ -Methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, 1 $\alpha$ -Isopropyl-1,3,4,5,6,11b-Hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-ol, 1 $\alpha$ -Ethyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-ol, (1 $\alpha$ -Ethyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, 1-Methyl-1 $\alpha$ ,3,4,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (1-Hydroxymethyl-1,3,4,5,6,11b-hexahydro-2H-11-oxa-4a-aza-

benzo[a]fluoren-1-yl)-methanol, 1-Methoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (-)-1-Methoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (+)-1-Methoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, 1 $\alpha$ -Methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene-1-carboxylic acid ethyl ester, 1-Ethoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (1 $\alpha$ -Methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, (-)-(1 $\alpha$ -Methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, (+)-(1 $\alpha$ -Methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-yl)-methanol, 1 $\alpha$ -Ethyl -1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene-1-carboxylic methyl ester, 1-Methoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (-)-1-Methoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (+)-1-Methoxymethyl-1 $\alpha$ -methyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene, (1 $\alpha$ -Ethyl-1,3,4,5,6,11b $\alpha$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluorene-1-yl)-methanol, acetic acid 1 $\alpha$ -Methyl-1,3,4,5,6,11b $\beta$ -hexahydro-2H-11-oxa-4a-aza-benzo[a]fluoren-1-ylmethyl ester or (1 $\alpha$ -Methyl-1,2,3,4,6,7,12,12b $\alpha$ -octahydroindeno[2,1-a]quinolizin-1-yl)-methanol.

23. (Currently Amended) A compound of formula IB



wherein,

X is NR<sub>2</sub>;

R<sub>2</sub> is (C<sub>1</sub>-C<sub>6</sub>)alkyl;

Z is -CHR<sub>8</sub>-(CH<sub>2</sub>)<sub>n</sub>- or a single bond;

R<sub>1</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, CN, NO<sub>2</sub>, NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino or carboxyl;

R<sub>3</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-

C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or one of R<sub>3</sub> or R<sub>4</sub> and R<sub>6</sub> together form a bond between the ring atoms to which they are attached;

R<sub>4</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or R<sub>4</sub> and R<sub>5</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring substituted with 1 to 3 substituent(s) R<sub>9</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>6</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>6</sub> forms a bond between the ring atom to which it is attached and the ring atom to which R<sub>7</sub> is attached;

R<sub>7</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>8</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or, only when n is 0, R<sub>7</sub> and R<sub>8</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring unsubstituted or substituted with 1 to 3 substituent(s) R<sub>10</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>15</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or carboxyl;

R<sub>16</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>7</sub> and R<sub>8</sub> are attached to the carbon ring atoms, which are adjacent;

m is 0 to 2; and

n is 0 or 1,

or a pharmaceutically acceptable salt or ~~and~~ ester thereof, with the provisos, that


a) when m is 0 or R<sub>1</sub> is methoxy and R<sub>4</sub> is H or ethyl, then R<sub>3</sub> is not methoxy-CO;

b) the compound is not 12-Methyl-1,2,3,4,6,7,12,12b-octahydro-indolo[2,3-a]quinolizine; 1-Ethyl-12-methyl-1,2,3,4,6,7,12,12b-octahydro-indolo[2,3-a]quinolizine; 2,3-Diethyl-12-methyl-1,2,3,4,6,7,12,12b-octahydro-indolo[2,3-a]quinolizine; 12-Methyl-1,2,3,4,6,7,12,12b-octahydro-indolo[2,3-a]quinolizine-1-ol; 2-(1-Ethyl-12-methyl-1,2,3,4,6,7,12,12b-octahydro-indolo[2,3-a]quinolizine-1-yl)-ethanol; 11-Methyl-2,3,5,6,11,11b-hexahydro-1H-indolizino[8,7-b]indole; (11-Methyl-2,3,5,6,11,11b-hexahydro-1H-indolizino[8,7-b]indol-1-yl)-methanol, (1,11-Diethyl-2,3,5,6,11,11b-hexahydro-1H-indolizino[8,7-b]indol-1-yl)-methanol or 3-(1-ethyl-12-methyl-1,2,3,4,6,7,12,12b-octahydro-indolo[2,3-a]quinolizine-1-yl)-propionic acid methyl ester.

24. (Original) A compound according to claim 23, wherein R<sub>3</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl and R<sub>4</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl or hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl.

25. (Currently Amended) A compound according to claim 23 ~~any one of claims 23 or 24~~, wherein the compound is 1 $\alpha$ -Ethyl-12-methyl-1,2,3,4,6,7,12b $\beta$ -octahydro-indolo[2,3-a]quinolizin-1-ol or 1 $\alpha$ -Ethyl-12-ethyl-1,2,3,4,6,7,12b $\beta$ -octahydro-indolo[2,3-a]quinolizin-1-ol.

26. (Currently Amended) A compound of formula IC



wherein,

X is NR<sub>2</sub>;

R<sub>2</sub> is H;

Z is -CHR<sub>8</sub>-(CH<sub>2</sub>)<sub>n</sub>- or a single bond;

n is 0;

R<sub>1</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, CN, NO<sub>2</sub>, NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino or carboxyl;



R<sub>3</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or R<sub>3</sub> and R<sub>6</sub> together form a bond between the ring atoms to which they are attached;

R<sub>6</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>6</sub> forms a bond between the ring atom to which it is attached and the ring atom to which R<sub>7</sub> is attached;

R<sub>7</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>8</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or, only when n is 0, R<sub>7</sub> and R<sub>8</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring unsubstituted or substituted with 1 to 3 substituent(s) R<sub>10</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl,

carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>9</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>15</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or carboxyl;

R<sub>16</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>7</sub> and R<sub>8</sub> are attached to the carbon ring atoms, which are adjacent;

m is 0 to 2; and

r is 1 to 3;


or a pharmaceutically acceptable salt or and ester thereof, with the provisos, that the compound is not 10-methyl-5,7,7a,8,9,10,11,11a,11b,12-decahydro-6H-6a,12-diaza-indeno[1,2-a]fluorene; 3-hydroxy-1,2,3,4,4a,5,6,7,8,13,13b,13c-dodecahydro-6a,13-diaza-indeno[1,2-c]phenanthrene-4-carboxylic acid methyl ester; methyl-3-ethyl-1,2,3a,4,6,7,12b,12c-octahydro-3H,12H-indolo[2,3-g]cyclopent[a]indolizine-2-carboxylate;

methyl-1,2,3a,4,6,7,12b,12c-octahydro-3H,12H-indolo[2,3-*g*]cyclopent[*a*]indolizine-2-carboxylate or 12c-ethyl-1,3a,4,6,7,12b,12c-octahydro-cyclopent[1,2]indolizino[8,7-*b*]indol-3(2H)-one.

27. (Original) A compound according to claim 26, wherein *r* is 1 and  $R_3$  is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl or hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl.

28. (Currently Amended) A compound according to claim 26 ~~any one of claims 26 or 27~~, wherein the compound is 3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ ,13 $\alpha$ -decahydro-2H-6a,13-diaza-indeno[1,2-*c*]phenanthren-1-one, 1,2,3,4,5,6,7,8,13,13b-decahydro-6a,13-diaza-indeno[1,2-*c*]phenanthrene, acetic acid 1 $\alpha$ ,2,3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ ,13 $\alpha$ -dodecahydro-6a,13-diaza-indeno[1,2-*c*]phenanthren-1-yl ester or acetic acid 1 $\beta$ ,2,3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ ,13 $\alpha$ -dodecahydro-6a,13-diaza-indeno[1,2-*c*]phenanthren-1-yl ester.

29. (Currently Amended) A compound of formula ID



wherein,

X is  $\text{NR}_2$ ;

$\text{R}_2$  is H;

Z is  $-\text{CH}(\text{CH}_2)_n-$ ;

n is 0;

$\text{R}_1$  is hydroxy,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , halogen, halo $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy-CO-}$ , CN,  $\text{NO}_2$ ,  $\text{NH}_2$ , mono- or di $(\text{C}_1\text{-C}_6)\text{alkylamino}$  or carboxyl;

$\text{R}_3$  is H, hydroxy,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_2\text{-C}_6)\text{alkenyl}$ , hydroxy $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}(\text{C}_1\text{-C}_6)\text{alkyl}$ , hydroxy $(\text{C}_1\text{-C}_6)\text{alkoxy}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$ ,  $(\text{C}_3\text{-C}_7)\text{cycloalkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryl, aryl $(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryloxy, aryl $(\text{C}_1\text{-C}_6)\text{alkoxy}$ , aryloxy $(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryl $(\text{C}_1\text{-C}_6)\text{alkoxy}(\text{C}_1\text{-C}_6)\text{alkyl}$ , halo $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $\text{NH}_2$ , amino $(\text{C}_1\text{-C}_6)\text{alkyl}$ , mono- or di $(\text{C}_1\text{-C}_6)\text{alkylamino}$ , mono- or di $(\text{C}_1\text{-C}_6)\text{alkylamino}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkyl-CO-}$ ,  $(\text{C}_1\text{-C}_6)\text{alkyl-CO-O-}$ ,  $(\text{C}_1\text{-C}_6)\text{alkyl-CO-O}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy-CO-}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy-CO}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy-CO}(\text{C}_1\text{-C}_6)\text{alkoxy}(\text{C}_1\text{-C}_6)\text{alkyl}$ , carbamoyl, mono- or di $(\text{C}_1\text{-C}_6)\text{alkylcarbamoyl}$ , carboxyl or  $(\text{C}_1\text{-C}_6)\text{alkyl-S}(\text{C}_1\text{-C}_6)\text{alkyl}$ , wherein the said  $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$  or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ , halogen,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ ,  $\text{NH}_2$ , CN or  $\text{NO}_2$ , or one of  $\text{R}_3$  or  $\text{R}_4$  and  $\text{R}_6$  together form a bond between the ring atoms to which they are attached;

$\text{R}_4$  is H, hydroxy,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ , hydroxy $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$  or  $(\text{C}_1\text{-C}_6)\text{alkoxy}(\text{C}_1\text{-C}_6)\text{alkyl}$ ;

$\text{R}_5$  is H, hydroxy,  $(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_2\text{-C}_6)\text{alkenyl}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}$ ,  $(\text{C}_1\text{-C}_6)\text{alkoxy}(\text{C}_1\text{-C}_6)\text{alkyl}$ ,  $(\text{C}_3\text{-C}_7)\text{cycloalkyl}$ ,  $(\text{C}_3\text{-C}_7)\text{cycloalkyl}(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryl, aryl $(\text{C}_1\text{-C}_6)\text{alkyl}$ , aryloxy,

aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or R<sub>4</sub> and R<sub>5</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring substituted with 1 to 3 substituent(s) R<sub>9</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>6</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>6</sub> forms a bond between the ring atom to which it is attached and the ring atom to which R<sub>7</sub> is attached;

R<sub>10</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>15</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-

CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or carboxyl;

R<sub>16</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;


R<sub>7</sub> and R<sub>8</sub> are attached to the carbon ring atoms, which are adjacent;

m is 0 to 2; and

t is 0 to 3;

or a pharmaceutically acceptable salt or ~~and~~ ester thereof, with the provisos, that the compound is not 1,2,3,4,4a,5,6,11,11b,12,13,13a-dodecahydro-4b,11-diaza-indeno[2,1-a]phenanthrene; 1,2,3,4,4a,5,6,11,11b,12-decahydro-4b,11-diaza-indeno[2,1-a]phenanthrene; 9-methoxy-1,2,3,4,4a,5,6,11,11b,12-decahydro-4b,11-diaza-indeno[2,1-a]phenanthrene or 1-hydroxy-1,2,3,4,4a,5,6,11,11b,12,13,13a-dodecahydro-4b,11-diaza-indeno[2,1-a]phenanthrene-2-carboxylic acid methyl ester.

30. (Currently Amended) A compound of formula IE



wherein,

X is NR<sub>2</sub>;

R<sub>2</sub> is H;

Z is -CHR<sub>8</sub>-(CH<sub>2</sub>)<sub>n</sub>- or a single bond;

R<sub>1</sub> is hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, CN, NO<sub>2</sub>, NH<sub>2</sub>, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino or carboxyl;

R<sub>3</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or one of R<sub>3</sub> or R<sub>4</sub> and R<sub>6</sub> together form a bond between the ring atoms to which they are attached;

R<sub>4</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>5</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryloxy, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy, aryloxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-

(C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl, carboxyl or (C<sub>1</sub>-C<sub>6</sub>)alkyl-S-(C<sub>1</sub>-C<sub>6</sub>)alkyl, wherein the said (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl or aryl is unsubstituted or substituted with 1 or 2 substituents each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, NH<sub>2</sub>, CN or NO<sub>2</sub>, or R<sub>4</sub> and R<sub>5</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring substituted with 1 to 3 substituent(s) R<sub>9</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>6</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or R<sub>6</sub> forms a bond between the ring atom to which it is attached and the ring atom to which R<sub>7</sub> is attached;

R<sub>7</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>8</sub> is H, hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy or (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl or, only when n is 0, R<sub>7</sub> and R<sub>8</sub> form, together with the carbon ring atoms to which they are attached, a condensed five to seven membered saturated carbocyclic ring unsubstituted or substituted with 1 to 3 substituent(s) R<sub>10</sub> each independently being hydroxy, (C<sub>1</sub>-C<sub>6</sub>)alkyl, halogen, NH<sub>2</sub>, NO<sub>2</sub>, (C<sub>3</sub>-C<sub>7</sub>)cycloalkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl,



carboxyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or oxo;

R<sub>15</sub> is H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, halo(C<sub>1</sub>-C<sub>6</sub>)alkyl, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylamino(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkyl-CO-O-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy-CO-(C<sub>1</sub>-C<sub>6</sub>)alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, carbamoyl, mono- or di(C<sub>1</sub>-C<sub>6</sub>)alkylcarbamoyl or carboxyl;

R<sub>16</sub> is H or (C<sub>1</sub>-C<sub>6</sub>)alkyl;

R<sub>7</sub> and R<sub>8</sub> are attached to the carbon ring atoms, which are adjacent;

m is 0 to 2; and

n is 1,

or a pharmaceutically acceptable salt or and ester thereof, with the proviso, that the compound is not 2,3,4,5,7,8,13,13b-octahydro-2,3-diethyl-1H-azepino[1',2':1,2]pyrido[3,4-*b*]indole; acetic acid 2,3,4,5,7,8,13,13b-octahydro-1H-azepino[1',2':1,2]pyrido[3,4-*b*]indol-2-ylmethyl ester; 2,3,4,5,7,8,13,13b-octahydro-1H-azepino[1',2':1,2]pyrido[3,4-*b*]indole-2-[(phenylmethoxy)methyl] or 2,3,4,5,7,8,13,13b-octahydro-1H-azepino[1',2':1,2]pyrido[3,4-*b*]indole-4-ethyl-2-[(phenylmethoxy)methyl].

31. (Original) A compound according to claim 30, wherein the compound is 2,3,4,5,7,8,13,13b-octahydro-1H-azepino[1',2':1,2]pyrido[3,4-*b*]indole.

32. (Original) A compound which is 2β-Methoxy-1,2,3,4,6,7,12,12bα-octahydro-indolo[2,3-*a*]quinolizine, 2α-methoxy-1,2,3,4,6,7,12,12bα-octahydro-indolo[2,3-*a*]quinolizine,

1 $\alpha$ -Ethyl-2 $\alpha$ -methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1-ol, 1 $\alpha$ -Isopropyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1-ol, (-)-1 $\alpha$ -isopropyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-ol, (+)-1 $\alpha$ -isopropyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-ol, 1 $\beta$ -Isopropyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizine, (1 $\alpha$ -Isopropyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1-yl)-methanol, (1 $\alpha$ - *n*-Propyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1-yl)-methanol, 2-(1 $\alpha$ ,2,3,4,6,7,12,12b $\beta$ -Octahydro-indolo[2,3-*a*]quinolizin-1-yl)-butan-2-ol, 1-(1,2 $\alpha$ ,3,4,6,7,12,12b $\alpha$ -Octahydro-indolo[2,3-*a*]quinolizin-2-yl)-propan-1-ol, 2-(1 $\alpha$ ,2,3,4,6,7,12,12b $\beta$ -Octahydro-indolo[2,3-*a*]quinolizin-1-yl)-propan-2-ol, 1-*s*-Butyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-ol, 1-Cyclohexyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-ol, 9-Fluoro-1 $\alpha$ -isopropyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1-ol, (1 $\alpha$ -Methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-yl)-methanol, (-)-(1 $\alpha$ -Methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-yl)-methanol, (+)-(1 $\alpha$ -Methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-yl)-methanol, (1 $\alpha$ -Ethyl-1,4,6,7,12,12b $\beta$ -hexahydroindolo[2,3-*a*]quinolizin-1-yl)-methanol, 3 $\beta$ ,4 $\alpha$ -Dimethyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizine, (1,2 $\alpha$ ,3,4,6,7,12,12b $\alpha$ -Octahydroindolo[2,3-*a*]quinolizin-2-yl)-propan-2-ol, (1,2 $\alpha$ ,3,4,6,7,12,12b $\beta$ -Octahydroindolo[2,3-*a*]quinolizin-2-yl)-propan-2-ol, (2 $\alpha$ -Ethyl-1,2,3,4,6,7,12,12b $\alpha$ -octahydroindolo[2,3-*a*]quinolizin-2-yl)-methanol, (2 $\alpha$ -Ethyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-2-yl)-methanol, (1 $\alpha$ -Ethyl-1,2,3,4,6,7,12,12b $\beta$ -octahydroindolo[2,3-*a*]quinolizin-1-ylmethoxy)-acetic acid ethyl ester, 1-(2 $\alpha$ -ethyl-1,2,3,4,6,7,12,12b $\alpha$ -octahydro-indolo[2,3-*a*]quinolizin-2-yl)-ethanone, 1-(2 $\alpha$ -ethyl-1,2,3,4,6,7,12,12b $\alpha$ -octahydro-indolo[2,3-*a*]quinolizin-2-yl)-ethanol, 2-(2 $\alpha$ -ethyl-

1,2,3,4,6,7,12,12b $\alpha$ -octahydro-indolo[2,3-*a*]quinolizin-2-yl)-propan-2-ol, 2-(3-ethyl-1,2 $\alpha$ ,3 $\alpha$ ,4,6,7,12,12b $\alpha$ -octahydro-indolo[2,3-*a*]quinolizin-2-yl)-propan-2-ol, (3-ethyl-2-methyl-1 $\alpha$ ,2 $\beta$ ,3 $\beta$ ,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1-yl)-methanol, 3-ethyl-1,2-dimethyl-1 $\alpha$ ,2 $\beta$ ,3 $\beta$ ,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizine, 1,2-dimethyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizin-1 $\beta$ -ol, (1-ethyl-2-methyl-1 $\beta$ ,2 $\beta$ ,3 $\beta$ ,4,6,7,12,12b $\alpha$ -octahydro-indolo[2,3-*a*]quinolizin-3-yl)-methanol, 1- $\beta$ -Hydroxymethyl-1-methyl-1,2,3,4,6,7,12,12b $\beta$ -octahydro-indolo[2,3-*a*]quinolizine-6 $\beta$ -carboxylic acid methyl ester, 5,6,7,7a $\beta$ ,8,9,10,11,11a $\beta$ ,11b $\alpha$ -Decahydro-12-oxa-6a-aza-indeno[1,2-*a*]fluorene, 2,3,4,4a $\beta$ ,5,6,7,8,13b $\beta$ ,13c $\beta$ -Decahydro-1H-13-oxa-6a-aza-indeno[1,2-*c*]phenanthrene, 2,3,4,4a $\beta$ ,5,6,7,8,13b $\alpha$ ,13c $\beta$ -Decahydro-1H-13-oxa-6a-aza-indeno[1,2-*c*]phenanthrene, 2,3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ -decahydro-1H-6a,13-diaza-indeno[1,2-*c*]phenanthren-13c $\beta$ -ol, (-)-2,3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ -decahydro-1H-6a,13-diaza-indeno[1,2-*c*]phenanthren-13c $\beta$ -ol, (+)-2,3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ -decahydro-1H-6a,13-diaza-indeno[1,2-*c*]phenanthren-13c $\beta$ -ol, (2,3,4,4a $\beta$ ,5,6,7,8,13,13b $\beta$ -Decahydro-1H-6a,13-diaza-indeno[1,2-*c*]phenanthrenyl)-13c $\beta$ -methanol or 5,6,7,7a,11,11b,12-Decahydro-6a,12-diaza-indeno[1,2-*a*]fluoren-11a-ol.

33. (Currently Amended) A pharmaceutical composition comprising at least one compound according to claim 15 ~~any one of claims 15 to 32~~ and a pharmaceutically acceptable diluent, carrier and/or excipient.

34. (Canceled)

35. (Canceled)

36. (New) A pharmaceutical composition comprising at least one compound according to claim 23 and a pharmaceutically acceptable diluent, carrier and/or excipient.

37. (New) A pharmaceutical composition comprising at least one compound according to claim 26 and a pharmaceutically acceptable diluent, carrier and/or excipient.

38. (New) A pharmaceutical composition comprising at least one compound according to claim 29 and a pharmaceutically acceptable diluent, carrier and/or excipient.

39. (New) A pharmaceutical composition comprising at least one compound according to claim 30 and a pharmaceutically acceptable diluent, carrier and/or excipient.